



Short Communication

# Neuronal death signaling pathway: Involvement of Rag1 activation

Takuma Hayashi<sup>1\*</sup>, Takao Hirano<sup>2</sup>, Toshinori Murata<sup>2</sup>

<sup>1</sup>Department of Immunology and Infectious Disease, Shinshu University Graduate School of Medicine, Matsumoto, Nagano, Japan

<sup>2</sup>Department of Ophthalmology, Shinshu University Graduate School of Medicine, Matsumoto, Nagano, Japan

\*Corresponding author email: [takumah@shinshu-u.ac.jp](mailto:takumah@shinshu-u.ac.jp)

**How to cite this article:** Takuma Hayashi, Takao Hirano, Toshinori Murata. Neuronal death signaling pathway: Involvement of Rag1 activation. IAIM, 2014; 1(4): 100-104.

**Available online at [www.iaimjournal.com](http://www.iaimjournal.com)**

**Received on:** 20-11-2014

**Accepted on:** 09-12-2014

## Abstract

Although the transcription factor, nuclear factor- $\kappa$ B (NF- $\kappa$ B) is known to regulate cell death and survival, its precise role in cell death within the central nervous system (CNS) remains unknown. We previously reported that mice with a homozygous deficiency for NF- $\kappa$ Bp50 spontaneously developed optic neuropathy. We examined the expression and activation of pro-apoptotic factor(s), which mediate optic neuropathy in p50-deficient ( $p50^{-/-}$ ) mice. Recombination activating gene 1 (Rag1) is known to regulate the recombination of immunoglobulin V(D)J. Experiments with genetically engineered mice revealed the involvement of Rag1 expression in the apoptosis of Brn3a-positive retinal ganglion cells (RGCs), and also showed the specific effects of a p50-deficiency on the activation of *Rag1* gene transcription. Furthermore, a genetic analysis of murine neuronal stem-like cells clarified the biological significance of Rag1 in NMDA (N-methyl-D-aspartate)-induced neuronal apoptosis. The apoptotic regulating factors, Bax, and cleaved caspase 3, 8, and 9 were observed in HEK293 cells expressing the external molecule of Rag1, and a human histological examination revealed the expression of Rag1 in RGCs. Recent study indicated that Rag1 played a role in optic neuropathy as a pro-apoptotic candidate in  $p50^{-/-}$  mice. This result may lead to new therapeutic targets in optic neuropathy.

## Key words

Rag1, NF- $\kappa$ Bp50, Programmed cell death, RGCs, Optic neuropathy.



The intracellular pathways related to cell survival regulate neuronal physiology during embryonic development as well as the pathogenesis of various neurodegenerative disorders. The NF- $\kappa$ B pathway was discovered in 1986 as a transcription modulator of the light chain of B lymphocyte immunoglobulins [1]. Subsequent studies have shown that NF- $\kappa$ B is a ubiquitously expressed dimeric transcription factor involved in numerous cellular processes, such as inflammation, differentiation, apoptosis, and oncogenesis. NF- $\kappa$ B is a dimer composed of members of the Rel family, which includes RelA(p65), RelB, and c-Rel [2]. The NF- $\kappa$ B family, which is primarily composed of p50/p65 (RelA) heterodimers, has been detected in almost animal cell types and is involved in cellular responses to stimuli such as stress and cytokines [3]. NF- $\kappa$ B is sequestered in the cytoplasm of unstimulated cells by a class of inhibitors called I $\kappa$ Bs. The degradation of I $\kappa$ B allows NF- $\kappa$ B to enter the nucleus, in which it specifically initiates the expression of target genes. Accordingly, the impaired regulation of NF- $\kappa$ B has been linked to various diseases, including cancer, inflammatory disorders, and autoimmune diseases, as well as deficiencies in the processes of synaptic plasticity and memory [4]. The NF- $\kappa$ B family also plays important roles in nervous system development and pathology by influencing neuronal apoptosis, neurite outgrowth, and synaptic plasticity [5, 6]. However, the range of intercellular signals and transduction mechanisms that regulate NF- $\kappa$ B activity in neurons is broad and complex. Knockout mice have been extensively used to assess different gene components in the NF- $\kappa$ B pathway. For instance, *p50*<sup>-/-</sup> mice exhibited the age-related degeneration of neuronal and non-neuronal cells, and defective NF- $\kappa$ B activation resulted in apoptosis in the striatal neurons of a Huntington disease model [7]. Activated NF- $\kappa$ Bp65 has been suggested to glutamate-induced neurotoxicity, N-methyl-D-aspartate (NMDA)-

induced retinal neuronal cell death, retinal ischemia, and reperfusion injury in the CNS [8, 9]. We previously reported that the number of retinal ganglion cells (RGCs) in *p50*<sup>-/-</sup> mice was significantly lower than that in its parental mice, *p50*<sup>+/+</sup> mice, suggesting that these animals exhibited features resembling those of human glaucoma [10]. However, the precise role of NF- $\kappa$ B in cell death within the CNS remains controversial. Therefore, we searched for a new target related to NF- $\kappa$ B pathways in neurons.

Verkoczy L, et al. reported that NF- $\kappa$ B was relevant in the B-cell receptor-mediated regulation of recombination activating gene (Rag) locus transcription [11]. They suggested that immediately activated NF- $\kappa$ B pathways may facilitate quick antigen receptor-regulated changes in Rag expression, which is important for editing [11]. *Rag* genes encode two enzymes that play key roles in the adaptive immune system: both Rag1 and Rag2 mediate the recombination of V(D)J, a process that is essential for the maturation of B and T cells in the development and maturation of lymphocytes [12, 13]. Rags have been detected not only in the immune systems of mammals and amphibians, but also in their nervous systems; *Rag1* transcripts have been found in the murine CNS, particularly in areas of high neural density, such as the cerebellum and hippocampal formation [14, 15, 16]. Rag1 may function in neurons to site-specifically recombine elements of the neuronal genome or prevent detrimental alternations in the genomes of long-lived cells. Although the role of the Rag1 locus in the CNS is currently unclear, Rags are known to be regulated by NF- $\kappa$ B [11]. Based on the above-described findings, we focused on Rag1 as a novel candidate target related to NF- $\kappa$ B pathways in neurons using *p50*<sup>-/-</sup> mice as a model of optic nerve neuropathy. Since no studies have been published on the expression of Rag1 in the visual system, we first confirmed



the presence of Rag1, but not the Rag2 transcript in RGCs. The absence of *Rag1* in *p50*<sup>-/-</sup> mice resulted in a decrease in optic nerve neuropathy. In vertebrate embryonic development, the retina and optic nerve originate as outgrowths of the developing brain, and, thus, the retina is considered to be part of the CNS. Furthermore, three-dimensional cultures of mouse embryonic stem cell aggregates have demonstrated the autonomous formation of the optic cup, which develops into the outer and inner layers of the retina structure from brain balls [17]. Glutamate is a major excitatory neurotransmitter in vertical pathways through the retina, wherein RGCs first express the NMDA glutamate receptors that are typical in the brain [18]. Since the brain and retina have a close relationship in genesis and neurotransmission, it is plausible that Rag1, which has been detected in the hippocampus, is also expressed in the retina.

We assessed the precise role of Rag1 in the retina using experiments with *p50*<sup>-/-</sup> mice, which exhibit age-dependent decreases in RGCs. A lack of Rag1 in *p50*<sup>-/-</sup> mice diminished the loss of RGC, which was confirmed by several lines of evidence in the present study. These results promoted us to speculate that Rag1 may play a role in the programmed cell death of RGCs, which was accelerated by NF- $\kappa$ B, in *p50*<sup>-/-</sup> mice. We found that Rag1 was also localized in the nucleus of RGCs in 15-month-old *p50*<sup>-/-</sup> mice whose RGC number had already markedly decreased; therefore we proposed that Rag1 may specifically influence apoptotic signaling in the nucleus [19]. **(Fig. 1)**

Many questions still remain regarding the molecular mechanisms involved in Rag1 functions in the retina. However, Rag1 may play a role in RGCs that is entirely distinct from somatic recombination. The evidence for this lies in studies of the molecular structure of the

recombinase enzymes themselves; although both Rag1 and Rag2 share several roles, i.e., DNA cleavage and rearrangement of V(D)J recombination, only Rag1 contained the catalytic DNA-binding core of the recombinase [20]. These domains are known to be similar to the active site of several transposases and integrases [21]. Kelch motifs, which mediate the interactions of Rag2 and Rag1, have been observed in numerous proteins [22], and Rag1 may interact with an identified protein via a kelch motif in the retina. As in the mouse retina, we confirmed the localization of Rag1, but not Rag2, in RGCs in the human retina. A protein homology study revealed that the human Rag1 molecule was 90% homologous with its mouse ortholog, the catalytic domains, zinc-finger, recombinase, and RING-finger in the Rag1 molecule appeared to be conserved between species. These results indicated the physiological significance of Rag1 observed in mice may extend to the regulation of human RGC survival. We concluded that Rag1 may also be involved in the programmed cell death of RGCs in the human glaucomatous retina. Further studies on the role of Rag1 in RGCs are expected to contribute to the development of preventive and therapeutic treatments for human glaucoma.

## References

---

1. Sen R, Baltimore D. Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell*, 1986; 46: 705-716.
2. Huxford T, Malek S, Ghosh G. Structure and mechanism in NF-kappa B/I kappa B signaling. *Cold Spring Harbor symposia on quantitative biology*. 1999; 64: 533-540.
3. Sha WC, Liou HC, Tuomanen EI, Baltimore D. Targeted disruption of the



- p50 subunit of NF-kappa B leads to multifocal defects in immune responses. *Cell*, 1995; 80: 321-330.
4. Hoffmann A, Leung TH, Baltimore D. Genetic analysis of NF-kappaB/Rel transcription factors defines functional specificities. *EMBO J*, 2003; 22: 5530-5539.
  5. Baeuerle PA, Baltimore D. I kappa B: A specific inhibitor of the NF-kappa B transcription factor. *Science*, 1988; 242: 540-546.
  6. Mattson MP, Meffert MK. Roles for NF-kappaB in nerve cell survival, plasticity, and disease. *Cell Death and Differentiation*, 2006; 13: 852-860.
  7. Lu ZY, Yu SP, Wei JF, Wei L. Age-related neural degeneration in nuclear-factor kappaB p50 knockout mice. *Neuroscience*. 2006; 139: 965-978.
  8. Fan W, Cooper NG. Glutamate-induced NFkappaB activation in the retina, *Investigative Ophthalmology & Visual Science*, 2009; 50: 917-925.
  9. Grilli M, Pizzi M, Memo M, Spano P. Neuroprotection by aspirin and sodium salicylate through blockade of NF-kappaB activation, *Science*, 1996; 274: 1383-1385.
  10. Takahash, Y, Katai N, Murata T, Taniguchi SI, Hayashi T. Development of spontaneous optic neuropathy in NF-kappaBetap50-deficient mice: Requirement for NF-kappaBetap50 in ganglion cell survival, *Neuropathol Appl Neurobiol*, 2007; 33: 692-705.
  11. Verkozy L, Ait-Azzouzene D, Skog P, Martensson A, Lang J, Duong B, Nemazee D. A role for nuclear factor kappa B/rel transcription factors in the regulation of the recombinase activator genes, *Immunity*, 2005; 22: 519-31.
  12. Mombaerts P, Iacomini J, Johnson RS, Herrup K, Tonegawa S, Papaioannou VE. RAG-1-deficient mice have no mature B and T lymphocytes. *Cell*, 1992; 68: 869-877.
  13. Oettinger MA, Schatz DG, Gorka C, Baltimore D. RAG-1 and RAG-2, adjacent genes that synergistically activate V(D)J recombination. *Science*, 1990; 248: 1517-1523.
  14. Chun JJ, Schatz DG, Oettinger MA, Jaenisch R, Baltimore D. The recombination activating gene-1 (RAG-1) transcript is present in the murine central nervous system. *Cell*, 1991; 64: 189-200.
  15. Feng B, Bulchand S, Yaksi E, Friedrich RW, Jesuthasan S. The recombination activation gene 1 (Rag1) is expressed in a subset of zebrafish olfactory neurons but is not essential for axon targeting or amino acid detection, *BMC Neurosci*, 2005; 6: 46.
  16. Fang M, Yin Y, Chen H, Hu Z, Davies H, Ling S. Contribution of Rag1 to spatial memory ability in rats. *Behav Brain Res*, 2013; 236: 200-209.
  17. Eiraku M, Takata N, Ishibashi H, Kawada M, Sakakura E, Okuda S, Sekiguchi K, Adachi T, Sasai Y. Self-organizing optic-cup morphogenesis in three-dimensional culture. *Nature*, 2011; 472: 51-56.
  18. Massey SC, Miller RF. Glutamate receptors of ganglion cells in the rabbit retina: evidence for glutamate as a bipolar cell transmitter. *The Journal of physiology*, 1988; 405: 635-655.
  19. Hirano T, Murata T, Hayashi T. Physiological significance of Rag1 in neuronal death, especially optic neuropathy. *FEBS J*, 2014, Oct 14. doi:10.1111/febs.13109.
  20. Fugmann SD, Lee AI, Shockett PE, Villey IJ, Schatz DG. The RAG proteins and V(D)J recombination: complexes, ends,


and transposition. Annual Review of Immunology, 2000; 18: 495-527.

21. Zhou L, Mitra R, Atkinson PW, Hickman AB, Dyda F, Craig NL. Transposition of hAT elements links transposable elements and V(D)J recombination. Nature, 2004; 432: 995-1001.
22. Prag S, Adams JC. Molecular phylogeny

of the kelch-repeat superfamily reveals an expansion of BTB/kelch proteins in animals. BMC Bioinformatics, 2003; 4: 42.

**Fig. 1:** Signal cascade of cell death mediated by Rag1 in p50-deficient mouse. A recent study demonstrated that the binding site of the hetero dimer p50-RelA(p65) could also be occupied by the homo dimer p50-p50, and may function as a repressor to regulate the role of p50-RelA(p65) as a transcription factor essential for neuronal responses. In p50-deficient neuronal cells, the c-Rel-RelA(p65) hetero dimer markedly induced *Rag1* gene activation as a transcription factor. Rag1 may play a role in neuronal cell death signaling as a nuclear mediator. The cell death factors, Bax and cleaved caspase-3, 8, and 9, were also clearly detected in Rag1-expressing cells.

Hayashi et al. Figure 1



**Source of support:** Nil

**Conflict of interest:** None declared.