

Original Research Article

# Retrospective study of fine needle aspiration cytology of clinically palpable breast lump

Gunvanti B. Rathod<sup>1</sup>, Mohit Jain<sup>2\*</sup>, Denish Vachhani<sup>2</sup>, Aviral Chandra<sup>2</sup>, Mansi Balar<sup>2</sup>

<sup>1</sup>Assistant Professor, Pathology Department, GMERS Medical College, Himmatnagar, Gujarat, India

<sup>2</sup>PG Student, Pathology Department, SBKS MI & RC, Sumandeep Vidyapeeth, Vadodara, Gujarat, India

\*Corresponding author email: [mohit2508@yahoo.co.in](mailto:mohit2508@yahoo.co.in)

	International Archives of Integrated Medicine, Vol. 3, Issue 2, February, 2016. Copy right © 2016, IAIM, All Rights Reserved. Available online at <a href="http://iaimjournal.com/">http://iaimjournal.com/</a> ISSN: 2394-0026 (P)                      ISSN: 2394-0034 (O)	
	Received on: 25-01-2016	Accepted on: 08-02-2016
	Source of support: Nil	Conflict of interest: None declared.
<b>How to cite this article:</b> Rathod GB, Jain M, Vachhani D, Chandra A, Balar M. Retrospective study of fine needle aspiration cytology of clinically palpable breast lump. IAIM, 2016; 3(2): 69-73.		

## Abstract

**Introduction:** Breast carcinomas are one of the leading causes of cancer in women. Fine Needle Aspiration Cytology (FNAC) is one of the important tools for the pre-operative diagnosis of breast lesions.

**Materials and methods:** A total of 200 breast aspirates were studied. Histo-cytopathological correlations were obtained in all cases. All the aspirates were stained with Hematoxylin and Eosin (H and E) stain.

**Results:** Among 200 patients, 195 were females and 5 were males. Benign breast lesions were found in 120 cases; among which fibroadenoma (34%) was the commonest lesion which was observed. Malignancy was observed in 80 cases; among them, ductal carcinoma was the predominant.

**Conclusion:** We can conclude that FNA is widely accepted as a reliable technique in the initial evaluation of palpable breast lumps. It is simple, safe, cost-effective, minimally invasive, rapid and as sensitive as biopsy.

## Key words

FNAC, Fine needle aspiration cytology, Breast lump, Benign, Malignant.

## Introduction

In both developed and developing countries, breast lumps comprise a considerable amount of surgical cases in women. In outpatient departments, breast lump is one of the commonest presentations; majority of them are in women and benign. Differentiating benign lumps from malignant preoperatively for definite treatment is necessary [1, 2]. The triple test which includes physical breast examination, mammography and fine-needle aspiration (FNA) has proved a reliable tool for accurate diagnosis of palpable breast masses. Fine needle aspiration cytology has become an increasingly popular technique utilized in the diagnosis of palpable breast masses owing to its distinct advantages of being sensitive and specific, expedient, economical and safe. The purpose of this study was to evaluate our experiences with fine needle aspiration cytology in a series of patient and compare the diagnostic accuracy of fine needle aspiration cytology with postoperative histopathology.

## Material and methods

The two years retrospective study of fine needle aspiration cytology of 200 clinically palpable breast lump with histopathological correlation was carried out during November 2012 to October 2014. FNAC was performed at cytology clinic using a 22-gauge needle attached to a 10 ml syringe after explaining procedure to the patients and obtaining their oral consent for the same [3-9]. The area to be aspirated was cleaned with spirit before aspiration and multiple hits were made within the lesion, with sufficient negative pressure; the needle was removed and the pressure was applied to the area of aspiration to avoid bleeding or hematoma formation. The aspirated material was smeared on glass slide and stained [10-16]. The cytological diagnosis was classified in to 3 groups benign, suspicious and malignant. After this reporting all the patients were later subjected to open/excision biopsy and its histopathological confirmation. Later diagnostic accuracy of cytology reporting was

compared with that of histopathology. The data was collected and analysed.

## Results

This study documented the fact that benign breast lesions were the most common lesions in young females, among which the Fibroadenoma was the commonest one. The malignant lesions were common in fourth and fifth decades of life, among which infiltrating ductal carcinoma was the most common lesion (**Table – 1, 2**).

**Table – 1:** Age wise distribution of cases.

Age (Years)	Patients	
	No.	%
≤20	08	04
21-30	32	16
31-40	89	44.5
41-50	60	30
51-60	06	03
>60	05	2.5
Total	200	100

**Table – 2:** Frequency of FNA diagnosis of 200 breast lesions.

Diagnosis	No. of cases	%
<b>Benign breast lesions (n = 120)</b>		
Fibroadenoma	68	34
Lactating adenoma	03	1.5
Fibrocystic change	20	10
Fibroadenosis	05	2.5
Gynecomastia	07	3.5
Galactocele	06	03
Benign phylloides	05	2.5
Inflammatory breast disease	09	4.5
<b>Malignant breast lesions (n= 80)</b>		
Ductal carcinoma	43	21.5
Mucinous carcinoma	08	04
Papillary carcinoma	10	05
Medullary carcinoma	07	3.5
Metaplastic carcinoma	06	03
Metastasis	06	03
<b>Total</b>	<b>200</b>	<b>100</b>

## **Discussion**

---

The fine needle aspiration cytology is an important diagnostic adjunct in the management of patient with a breast lump [17]. Its distinct advantage is that it can be done during the outpatient visit without the need of the anesthesia, thus eliminating the cost of outpatient surgery. It also allows discussion with the patient of various treatment plans for malignant mass on the same visit. It greatly compliments the clinical and radiological examination and permits rapid diagnosis in more than 95% of the cases.

In our study, we had 120 benign lesions (60%), fibroadenoma being the most common benign lesion (56.67%) that presents for needle aspiration. This has been confirmed in other series also. The fibroadenoma has been considered a significant cause for the false positive diagnosis. The overall activity of the epithelial cell in this tumor is probably the reason. We had no cases of false positive reports in our study. Ductal carcinoma is one of the most common malignancies (53.75%) among women in our study. The breast lump is usually discovered by the patient. The typical carcinoma presents a gritty resistance to the fine needle. The aspirate is usually copious and blood stained.

Another common benign breast lesion we encountered was fibrocystic change ( $N = 20$ ) with maximum patients between 21-40 years. Though hormones play a role in its development exact pathogenesis remains obscure [18]. Fibrocystic change is not a specific cytological diagnosis. Cytology samples must be evaluated in the context of clinical and mammography findings. Some of these lesions simulate carcinoma clinically, radiologically, and microscopically [19]. More than 90% of the fibrocystic changes were non-proliferative and FNA smears showed many macrophages, apocrine cells with or without scanty chronic inflammatory cells.

However, one case which was misinterpreted as a benign cystic lesion by FNAC, was later on

diagnosed as a malignant phyllodes tumour on doing a histopathological examination. This might be due to inadequate sampling, because of the cystic nature of lesion. So, in case of cystic lesions, it is better to re-aspirate the lesion from the solid area after evacuation of cyst or image guided FNA should be performed to locate solid area. It is always necessary to correlate the FNAC findings with clinical diagnoses and mammograms and to go for core biopsies whenever they are needed, to avoid misdiagnoses. The false negative rate varies from 1-8% in different studies [20-23].

In the present study, all the 80 cytologically diagnosed malignant cases were confirmed as malignant on subsequent histopathological examinations. So, in our study, a 100% cyto-histopathological correlation was observed for malignant lesions. Zhang Qin et al. [22], AZ Mohammed et al. [23], Tiwari M [20] had also observed the same results in their studies.

Apart from the high accuracy rate of fine needle aspiration cytology, this technique is quite attractive because of its rapidity of execution and interpretation. Some have raised questions about the possible dangers of cell implantation from the needle aspiration. These rare reports have largely resulted from the use of larger cutting needle (18 gauge) rather than fine needles (22 gauge). With this fine needle technique, there is essentially no danger of implantation with breast aspiration [24]. Franzen and Zajicek in a review of 3479 consecutive breast aspirates found no evidence of seeding along the needle tract [25]. This is not surprising as the needle tract is invariably removed with definitive surgery.

## **Conclusion**

---

We can conclude that fine needle aspiration (FNA) cytology is a safe, cost-effective, and reliable technique for preoperative evaluation of palpable breast lumps. FNA features are more informative when combined with clinical and radiological findings. Clinical breast examination and mammography screening in female subjects

should be encouraged in developing countries from the third decade onward for early detection of breast carcinoma.

## References

1. Vaidyanathan L, Barnard K, Elnicki DM. Benign breast disease: When to treat, when to reassure, when to refer. *Cleve Clin J Med.*, 2002; 69: 425-32.
2. Guray M, Sahin AA. Benign breast diseases: Classification, diagnosis, and management. *Oncologist*, 2006; 11: 435-49.
3. Gunvanti Rathod, Pragnesh Parmar. Fine needle aspiration cytology of swellings of head and neck region. *Indian Journal of Medical Sciences*, 2012; 66: 49-54.
4. Gunvanti Rathod, Sangita Rathod, Pragnesh Parmar, Ashish Parikh. Diagnostic efficacy of fine needle aspiration cytology in cervical lymphadenopathy – A one year study. *International Journal of Medical and Pharmaceutical Sciences*, 2014; 4(5): 1-8.
5. Rathod GB, Ghadiya V, Shinde P, Tandan RK. Pleomorphic sarcoma in 60 years old male – A case report. *International Journal of Current Microbiology and Applied Sciences*, 2014; 3(8): 510-517.
6. Gunvanti Rathod, Pragnesh Parmar, Sangita Rathod, Ashish Parikh. Suprascapular malignant fibrous histiocytoma – A case report. *Discovery*, 2014, 12(31): 50-53.
7. Rathod GB, Goyal R, Bhimani RK, Goswami SS. Metaplastic carcinoma of breast in 65 years old female - A case report. *Medical Science*, 2014; 10(39): 77-81.
8. Disha Singla, Gunvanti Rathod. Cytodiagnosis of renal cell carcinoma – A case report. *IAIM*, 2015; 2(2): 133-137.
9. Mobeen Alwani, Gunvanti B. Rathod. Diagnosis of anaplastic thyroid carcinoma on fine needle aspiration cytology - A rare case report. *IAIM*, 2015; 2(3): 183-187.
10. Annie Jain, Gunvanti Rathod. Oncocytoma of parotid gland: A rare case report. *IAIM*, 2015; 2(4): 166-169.
11. Nupur Singla, Gunvanti Rathod, Disha Singla. Adenoid cystic carcinoma of the parotid gland - A case report and review of literature. *IAIM*, 2015; 2(4): 182-186.
12. Anchal Bhola, Gunvanti Rathod, RK Tandan. Cystic metastatic squamous cell carcinoma - A case report. *IAIM*, 2015; 2(5): 195-199.
13. Rathod GB, Jain A. Role of FNAC in diagnosis of gouty tophi - A case report. *IAIM*, 2015; 2(7): 137-140.
14. TH Kalidas Singh, Gunvanti B. Rathod. Diagnosis of fat necrosis on FNAC - A case report. *IAIM*, 2015; 2(6): 236-239.
15. Rathod GB, Rai P. Audit of repeat fine needle aspiration in cytopathology laboratory. *IAIM*, 2015; 2(9): 20-25.
16. Rathod GB, Rai P, Rai S. A prospective study of ultrasonographic and FNAC correlation of thyroid pathology. *IAIM*, 2015; 2(11): 46-51.
17. Hussain MT. Comparison of fine needle aspiration cytology with excision biopsy of breast lump. *J Coll Physicians Surg Pak.*, 2005; 15(4): 211-214.
18. Rosai J. *Rosai and Ackerman's Surgical Pathology*, Vol. 2, 9<sup>th</sup> edition, Missouri: Elsevier Publishers; 2004, p. 1779.
19. Marshall LM, Hunter DJ, Connolly JL, Schnitt SJ, Byrne C, London SJ, et al. Risk of breast cancer associated with atypical hyperplasia of lobular and ductal types. *Cancer Epidemiol Biomarkers Prev.*, 1997; 6: 297-301.
20. Tiwari M. Role of FNAC in diagnosis of breast lumps. *Kathmandu University Medical Journal*, 2007; 5: 215-17.
21. O'Neil S, Castelli M, Gattuso P, Kluskens L, Madsen K, Aranha G. Fine-needle aspiration of 697 palpable breast lesions with histopathologic correlation. *Surgery*, 1997; 122(4): 824-28.

22. Zhang Qin, Nie Shigui, Chen Yuhua, Zhou Limei. Fine Needle Aspiration Cytology of Breast Lesions: Analysis of 323 Cases. *The Chinese-German Journal of Clinical Oncology*, 2004; 3(3): 172-74.
23. Mohammad AZ, Edino ST, Ochicha O, Alhassan SU. Value of fine needle aspiration biopsy in preoperative diagnosis of palpable breast lumps in resource-poor countries: a Nigerian experience. *Annals of African Medicine*, 2005; 4(1): 19-22.
24. Michael Shabot. Aspiration Cytology is Superior to Tru-cut Needle Biopsy in Establishing the Diagnosis of Clinically Suspicious Breast Masses. *Annals of Surgery*, 1982; 196: 122-126.
25. Franzen S, Zajoak J. Aspiration biopsy in diagnosis of palpable lesions of the breast. *Acta Radiol.*, 1968; 7: 241-262.