

Original Research Article

Serial CT Scan following Traumatic Brain Injury: Its Significance and Analysis

I.D. Chaurasia*, Anshul Siroliya, Manish Parihar

Department of Surgery, Gandhi Medical College and Hamidia Hospital, Bhopal, Madhya Pradesh, India

*Corresponding author email: chaurasiaid@gmail.com

	International Archives of Integrated Medicine, Vol. 5, Issue 5, May, 2018. Copy right © 2018, IAIM, All Rights Reserved. Available online at http://iaimjournal.com/	
	ISSN: 2394-0026 (P)	ISSN: 2394-0034 (O)
	Received on: 16-04-2018	Accepted on: 22-04-2018
	Source of support: Nil	Conflict of interest: None declared.
How to cite this article: I.D. Chaurasia, Anshul Siroliya, Manish Parihar. Serial CT Scan following Traumatic Brain Injury: Its Significance and Analysis. IAIM, 2018; 5(5): 1-7.		

Abstract

Introduction: CT plays a key role in accurate diagnosis and important diagnostic tool in the management of traumatic brain injury.

Materials and methods: Present study was under taken in the Department of Surgery Gandhi Medical College and Hamidia Hospital, Bhopal in the duration of July 2015 to July 2016. Total 105 patients (M: 80, 76.1% and F: 25; 23.80%) with traumatic head injury were admitted and managed and were include in this study. They were followed by serial CT scan and analyzed.

Results: Commonest cause of brain injury was road traffic accidents (75 cases 70%). Altered sensorium was the most common symptom in 88 cases (88.0%), followed by vomiting in 49 cases (75.38%). Contusions were the most common radiological findings in 80.95% (85 cases) followed by extradural haematoma in 18.46% and pneumocephalus in 16.92%. Out of these the most common location for contusion was frontal (14 cases), followed by parietal (12 cases) and then bilateral contusions in 10 cases. Repeat scan showed increase in volume of contusion in 29 cases (44.61%), no change in 26 cases (40.0%) and decrease in 10 cases. The increase was mainly due to edema in 18 cases (27.69%) and due to actual increase in contusion in only 11 cases. Only 4 cases needed operative intervention.

Conclusions: Sequential serial CT scan results in the management changes and has role in the management even in the patients without any clinical deterioration in early stages.

Key words

Serial CT scan, Traumatic Brain injury, GCS.

Introduction

Traumatic brain injuries are one of the most commonly encountered life threatening trauma emergency. The treatment of head injuries depends on the type of injury and the patient's condition. To assess the severity of a head injury, a physician may perform a physical and neurologic exam and imaging tests. Computed tomography represents a major breakthrough in the investigation of cases of head injuries. In management of craniocerebral trauma patients it has become the single, sufficient, and necessary radiographic test. Although it is used widely in trauma the indications for repeat scan are not well defined. In patients managed conservatively the role of repeat scan has been a subject of debate [1]. The extent of information the second or repeat scan provides than the initial scan or control scan varies from study to study. Different studies have reported conflicting conclusions on the advantage and disadvantage of doing routine CT scan. Since majority of the relevant pathological changes develop within the first twenty four hours a pathological categorization using the initial scan has also been proposed for prognostic purpose.

Materials and methods

Present study was conducted in Department of Surgery Gandhi Medical College and Hamidia Hospital, Bhopal. It was a prospective study that included 105 cases of traumatic brain injury admitted and initially managed conservatively at Hamidia hospital, Bhopal from November 2015 to July 2016. Repeat CT scan head was performed on a routine basis at 24 hours of admission unless the patient deteriorated in which case it was done earlier. The sex ratio, age, mode of injury, presenting symptoms will be studied. All the pre and post repeat scan radiological findings was recorded with respect to appearance of new lesions, initial volume, and change in lesion or edema. The initial Glasgow Coma Score (GCS) at admission and the GCS at time of repeat scan was noted. The change in operative management and the advantage of repeat scan in terms of prognosis of head injury

patients when used routinely was assessed. As per our methodology all those cases with significant mass lesion that required surgery, or cases that had Intracranial Pressure Monitoring (ICP) insertion and those cases which were admitted more than 12 hours of injury were excluded from this study. Repeat scan was done as a routine basis after 24 hours of admission irrespective of the GCS score (**Figure – 1, 2**).

Figure – 1: CT scan within 6 hours of admission showing small acute subdural hematoma.





Figure – 2: CT scan after 24 hours shows Right temporo-parietal contusion with edema.



Results

The total number of cases included in this study was 105 with males 80 patients, 70.76% more common than females (25 patients, 23.80%). The majority of patients were between age group's 21-30 years 38.09% and 31-40 years 24.76% respectively (**Table - 1**). The age group ranged from 10 years to 70 years.

Table - 1: Age Groups.

Age Groups	N=105	%
10-20	10	9.52 %
21-30	40	38.09 %
31-40	26	24.76 %
41-50	14	13.33 %
51-60	09	8.57 %
61-70	06	5.71 %
Range from 10 years to 70 years		

In our study road traffic accidents cases were 66.66 % (70 cases) of head injury. Assault was the next common cause 19.04 % (20 cases) (**Table - 2**).

Table - 2: Mode of injury.

Mode of Injury	Number	%
R.T.A.	70	66.66
Assault	20	19.04
Fall from Height	15	14.28

Altered sensorium 85.71% (90 cases), vomiting 80.95% (80 cases) and headache 61.90% (65 cases) were the most common modes of presentation in head injury patients. Early seizures defined as within one week of injury was present in 09 cases (13.84%) (**Table - 3**).

Contusions, including bilaterally and counter coup types, were the most common radiologic findings in 74.28 % (78 cases) followed by pneumocephalus 18.09 % (19 cases), depressed fracture (17 cases) 16.19%, Subdural hematoma 15.23% (16 cases) and extradural hematoma in 5.71%. Out of these the most common location for contusion was frontal (28 cases 35.89%), followed by parietal (26 cases 33.33 %) and then

bilateral contusions (18 cases) (**Table - 4**). Repeat scan showed increase in volume of contusion in 29 cases (44.61%), no change in 26 cases (40%) and decrease in 10 cases (15.83%). The increase was mainly due to edema in 18 cases (27.69%) and due to actual increase in contusion in only 11 cases (**Table - 5**). There was no increase in size of the extradural hematoma on the repeat scans (**Table - 6**).

Table - 3: Symptoms present.

Symptoms	No. of Cases	%
Altered Sensorium	90	85.71
Vomiting	85	80.95
Headache	65	61.90
E.N.T. bleed	28	26.66
Black Eye	42	40.0
Seizure	14	13.33
Associated Injury	43	40.95

Table - 4: Initial CT Scan Finding.

Finding	No.	%
Contusion	78	74.28
EDH	6	5.71
Pneumocephalus	19	18.09
SAH	12	11.42
Subdural Haematoma	16	15.23
Fracture Depressed/ Elevated	17	16.19
Intra Cerebral Haematoma	07	6.66

The admission GCS was 13-15 in 28 cases (43.07%), 8-12 in 33 cases (50.76%) and only 4 cases with 6.0% had score less than 7. Thus the majority of cases were in the moderate group. Repeat score during the scan at 48 hours showed change from 28 (43.07%) to 35 (53.84%) cases for score of mild head injury and decrease from 33 (50.76%) to 29 (44.61%) for moderate head injury. This showed clinical improvement to score of 13-15 in 10% of cases within 24 hours of admission (**Table - 7**).

Change in the management plan by operative intervention was needed in 17 cases (6.15%). Of these 4 cases two had a GCS of 15, one had score of 10 and one had score of 12 on initial

admission. Three out of these four had associated drop in GCS and the other had increase in contusion volume requiring intervention. Considering the location of the lesions 3 of them were in the frontal lobe and one in temporal lobe. All of them improved after surgery. **Table - 8** shows that most of the patient comes early to

hospital and their CT scan done early. Most of the head injury patient underwent CT scan head within 3-6 hours of head injury. Patients who had repeat CT scan head after 12 hours of head injury were excluded. Maximum changes occur in repeat CT scan of those patients who had CT scan early after head injury.

Table - 5: Admission and repeat scan findings.

Findings	No. N=105	Initial Volume (mean)	Repeat Volume (mean)	Surgical Intervention
Contusion	78			
Frontal	28	12	14	5 cases
Parietal	26	9	10	
Temporal	14	9	12	12case
Occipital	01	4	4	
Mixed: Fronto-Temporal	11	5	5	
Bilateral Contusions	18	9	11	
EDH	5	5	6	
Pneumocephalus	19	12	10	

Table - 6: Repeat Scan Findings.

Result	No.	%
No Change	43	40.95
Increase of Haematoma/contusion	47	44.76
Decrease	16	15.23
Increase Edema/mass effect	30	28.57
Appearance of new lesions	17	16.19

Table - 7: GCS Change with regards to repeat scan and outcome.

GCS	Admission GCS	Repeat GCS in 48 Hours	GCS Improvement
13-15	31 (29.52%)	43 (40.95%)	18%
8-12	47 (44.76.%)	43 (40.95%)	
3-7	7 (6.66%)	3 (2.85%)	

Table - 8: Time between injury and CT scan and changes in repeat scan.

Time between Injury and CT Scan	No. of Patients	No. and % of Patients in Which Changes Occurs in Repeat CT Scan	
		No.	%
0-3 hours	8	9	75
3-6 hours	27	30	70.37
6-9 hours	2	16	50
9-12 hours	10	7	40

Discussion

CT scan is one of the important diagnostic tool in evaluation of head injury. During the acute period, it permits the rapid, safe detection and precise localization of intracranial hematomas, brain contusion, edema and foreign bodies and serial CT scan aid in the diagnosis of subsequent complications. The exact time and the result of repeat scans yet to be determine.

Many of them have been studied on severe or blunt head injury. The old study to record repeat scan was in the study by French and Dublin in 1977 in which out of their series of 1000 head injury, cases 103 had repeat scans 31% of them undergoing surgical intervention.[2] One of the earliest study dedicated to repeat scan by Roberson et al. in 1979 concluded that repeat scan could be done on first post craniotomy day, on third day if no improvement up to 2-3 weeks and 3 months. They saying that repeat scan were of value in identifying post-traumatic hydrocephalus, delayed intracerebral hematomas, and intraventricular hemorrhage [3].

In study of Velmahos in Boston they reviewed 179 patients of minimal head injury and repeat CT scan were done in all patient, of them 37 (21%) showed signs of injury evolution on repeat CT scan and 7 (4%) require intervention. All 7 (4%) had clinical deterioration preceding CT scan. With this study they concluded repeat CT scan is unnecessary in patients with minimal head injury. Clinical examination identifies accurately the few who will show significant evaluation and require intervention [4].

In study by S. B. Dharap in 2004 in Mumbai India, 175 patients of head injury were reviewed and repeat CT scan was done and there was no standard protocol for ordering the repeat CT scans. CT scan was repeated in 53 (30%) patients. The clinical indication for the repeat CT scan in their study grouped into three (a) clinical deterioration (b) failure of

improvement (c) as a follow-up scan. Nine underwent surgical intervention based on the repeat CT scan finding; in others the repeat CT scan finding did not alter the management. They were concluded that in patients of head injury who are improving follow up repeat CT scan is unnecessary and unlikely to yield any further information necessitating change in treatment [5]. In study by Y. B. Roka in 2008 in Nepal, 71 cases of head injury were reviewed and underwent repeat CT scan head after 48 hours after admission. Repeat scan showed increase in volume of contusion in 31 cases, no change in 28 cases and decrease in 12 cases. Only five cases needed surgery. They were concluded that routine use of repeat scan in mild to moderate head injury has no role unless there is clinical deterioration.

Since that study the literature is also full of papers that support or disagree with serial CT scan in head injury patients [9-12]. Those of them which support have called it standard practice to repeat scans at 1, 3, 7 days and 1 month or on days 3 and 7 [1]. Other studies have concluded that repeat scan should be done at 12 hours if the first scan was done within 3 hours of injury the reason being that a significant proportion of cases, especially if unconscious may be harboring intracranial lesions [6]. Good outcome was found in cases who did not have new lesions on repeat scan [7].

Various indications for repeat CT scan have been proposed.

Frankhauser, et al. in 1983 proposed the following [8]:

- After 2-3 days if the first CT scan shows a fracture with an overlying collection of blood too small for surgical intervention.
- In all cases of secondary deterioration.
- Where there is a secondary elevation of intracranial pressure.
- Where there is no clinical improvement.

- In cases where there is persistent elevation of ICP after evacuation of intracranial hemorrhage.
- After 12 to 24 hours for all paralyzed and ventilated patients.

Bucci, et al. in 1986 suggested that poly trauma patients must be rescanned after correction of hypovolemia even if the initial CT scan was normal [9].

In addition a repeat scan within 24 hours was also suggested to detect a delayed contra lateral hematoma after surgical decompression.

Lately the indication for repeat CT have been simplified, Gentleman, et al. in 1989 mentioned they repeat CT is indicated when [10]

- Mean ICP becomes elevated above 30 mmHg.
- There is clinical deterioration.
- There is failure to improve

A similar protocol was advocated by Handley in 1991. In this study we took 24 hours for repeat scan as the maximal changes occur within the same period and any advantage thereby provided by repeat scan would be most beneficial. The changes in repeat scan in this study were present in 44.61%, which was mainly due to edema (27.69%). Study by Roka, et al. showed 43.6% new lesion development.

Contusions were the most common findings and the frontal lobe as the most common site. This would reflect direct injury. Repeat scan findings that needed surgery was found in only 6.15% of cases in this study and the majority of them were located in the frontal lobe. This is in accordance with the study by Patel et al. which show that in conservatively managed patient's frontal intra parenchymal hematoma were more prone to failure [11].

A similar study by Sifri, et al. on patients with minimal head injury with a bleed showed that out of 161 cases in their study only 4.0% needed

intervention and none of those with repeat scan needed intervention. The Glasgow outcome scale in this study was favorable in 98.0% of cases. With this they concluded that repeat scan would have no benefit and thereby suggested multicenter trial to further validate it [12]. Another study by Brown, et al., who did a prospective study of 100 patients who underwent repeat scan showed that of the 68.0% who underwent repeat scan 90.0% was done on routine basis and only 10.0% due to a change in ICP, GCS, both pupil changes or new onset of headache.. In study by Roka of 71 patients 31 patients showed increased hematoma and only 5 underwent surgery. Sequential CT scan should be performed as it may decide or change the treatment plan of the patient.

Conclusion

We conclude that serial sequential CT scan should be performed in traumatic brain injury patients when there is clinical deterioration, neurological deficit, papillary changes and in old age and also change in motor system as well change in the GCS. Repeat CT scan gives the vital information and has got a significant role in management of traumatic brain injury.

References

1. Roka YB, Kumar P, Bista P, Sharma GR, Adhikari D, Khadka NK, et al. Role of repeat CT scan in initially inoperable cases of traumatic head injury. Nepal Med Coll J., 2008; 10: 225-9.
2. French BN, Dublin AB. The value of CT scanning in the management of 1000 consecutive cases of head injuries. Surg Neurol., 1977; 7: 171-83.
3. Roberson FC, Kishore PR, Miller JD, Lipper MH, Becker DP. The value of serial computerized tomography in the management of severe head injury. Surg Neurol., 1979 Aug; 12(2): 161-7.
4. Velmahos G, Gervarini A, Petrovick L, Dorer DJ, Doran ME, Spaniolas K, et al. Routine repeat head CT for minimal

- head injury in unconscious. *J Trauma*, 2006; 60: 494-501.
5. Dharap SB, Akhandekar A, Pandey A, Sharma AK. Repeat CT scan in closed head injury. *Injury*, 2005; 36(3): 412-6.
 6. Servadei F, Nanni A, Nasi MT, Zappi D, Vergoni G, Giuliani G, et al. Evolving brain lesions in the first 12 hours after head injury: analysis of 37 comatose patients. *Neurosurgery*, 1995; 5: 899-907.
 7. Kobayashi S, Nakazawa S, Otsuka T. Clinical value of serial computed tomography with severe head injury. *Surg Neurol.*, 1983; 30: 25-9.
 8. Frankhauser H, Kener M. Delayed development of extradural haematoma. *Acta Neurochir.*, 1982; 60: 29-35.
 9. Bucci MN, Phillips TW, McGillicuddy JE. Delayed epidural hemorrhage in hypotensive multiple trauma patients. *Neurosurgery*, 1986 Jul; 19(1): 65-8.
 10. Gentleman D, Nath F, Macphersan P. Diagnosis and management of delayed traumatic intracerebral haematomas. *Br J Neurosurg.*, 1982; 3(3): 367-72.
 11. Patel NY, Hoyt DB, Nakaji P, Marshall L, Holbrook T, Coimbra R, et al. Traumatic brain injury: patterns of failure of non-operative management. *J Trauma*, 2000; 48: 367-74.
 12. Sifri ZC, Homnick AT, Vaynman A, Lavery R, Liao W, Mohr A, et al. A prospective evaluation of the value of repeat cranial computed tomography in patients with minimal head injury and an intracranial bleed. *J Trauma*, 2006; 61: 862-7.