


Original Research Article

Study of Prognostic Significance of Anti Cyclic Citrullinated Peptide Antibodies in Patients with Rheumatoid Arthritis

Haresh Panchal*

Assistant Professor, Department of Medicine, BJ Medical College, Civil Hospital, Ahmedabad, Gujarat, India

*Corresponding author email: hareshpnchl@gmail.com

	International Archives of Integrated Medicine, Vol. 6, Issue 5, May, 2019. Copy right © 2019, IAIM, All Rights Reserved. Available online at http://iaimjournal.com/	
	ISSN: 2394-0026 (P)	ISSN: 2394-0034 (O)
	Received on: 12-04-2019	Accepted on: 17-04-2019
	Source of support: Nil	Conflict of interest: None declared.
How to cite this article: Haresh Panchal. Study of Prognostic Significance of Anti Cyclic Citrullinated Peptide Antibodies in Patients with Rheumatoid Arthritis. IAIM, 2019; 6(5): 1-4.		

Abstract

Background: Rheumatoid arthritis is a chronic autoimmune inflammatory disease with articular and extra-articular manifestation that affects 0.5 to 1 % of total population.

Aim and Objectives: To correlate Rheumatoid factor (RF) and Anti-CCP antibody in RA, to evaluate prognostic value of RF and Anti-CCP antibody in RA.

Materials and methods: Retrospective study of total 50 patients admitted to our hospital from January 2018 to December 2018 was done. All patients were diagnosed as rheumatoid arthritis as per diagnostic criteria of American College of Rheumatology. All patients had symptom duration of at least one Year. Anti-CCP and IgM-RF were evaluated in all patients. Disease activity score 28 was calculated in all patients. Radiological Damage was assessed by Larsen Score.

Results: Anti-CCP and RF were significantly correlated with each other and both were seen as significant independent predictors of radiological outcomes (p value 0.01 and <0.05 respectively). Combination of these two had highest risk for erosive joint damage.

Conclusion: RA is more common in female. Anti-CCP antibody and RF both in combination were associated with higher probabilities of erosive disease.

Key words

Rheumatoid Factor(RF), Anti-CCP Antibody (Anti-ccp), ESR, CRP, Rheumatoid arthritis (RA), Larsen Score, DAS28, DMARD, Autoimmune, Intra-Extra articular.

Introduction

Rheumatoid arthritis is a chronic autoimmune inflammatory disease with articular and extra-articular manifestation that affects 0.5 to 1% of total population [1, 2]. Diagnosis of RA depends on clinical symptoms, lab investigation and imaging [3]. Joint damage accounts for a considerable part of the disability caused by RA [4]. Most effective way to manage RA patients is early diagnosis and timely treatment with disease modifying anti rheumatic drug which prevent the exacerbation of disease and improves outcome [5, 6]. Availability of better prognostic marker would make it possible to select predictably severe cases for aggressive therapy at early stage [7, 8].

Antibodies against cyclic citrullinated peptide (anti-CCP) are a new and highly specific maker for RA [9, 10]. Anti-CCP antibodies are now considered as an important serological marker for the diagnosis of RA and as a possible prognostic marker for development of erosive disease [11, 12].

Aims and objectives

- To correlate RF with Anti-CCP antibody
- To determine predictive value of anti-CCP and RF for radio-logical damage in patients with RA.

Materials and methods

Retrospective study of 50 patients who were diagnosed as rheumatoid arthritis as per diagnostic criteria of American college of rheumatology. All patients had symptom duration of minimum 1 year. Clinical evaluation of disease was based on number of swollen joints, number of tender joints, and morning stiffness duration. Disease activity was assessed by 28 joint disease activity score (DAS28).

Radiographic measurement: Standardized Radiographs of hands were performed to calculate the Larsen score. In each case, 22 joints were scored: all metacarpophalangeal joints (=10), all proximal interphalangeal joints (=8),

both first interphalangeal joints (=2) and wrists (=2) in the hands, Each joint grading in all 22 joints, ranging from 0 to 110. Joints damage was defined if Larsen score was 10 or higher, otherwise it was not present.

RF and antibodies to cyclic citrullinated peptide were analyzed. Anti-CCP was considered positive if titre was greater than 5 IU/ml. IgM-RF was determined positive if result of >20 IU/ml. ESR (erythrocyte sedimentation rate) and CRP (C-reactive protein) was measured in the all patients.

Results

Mean age of patients was 44 years and mean duration of disease was 9 years. 40 (80%) of all 50 patients were RF positive while 10 (20%) were negative. 37 (74%) of all patients were positive for anti-CCP while 13 (26%) were negative. 30 patients (60%) were positive for both RF and anti-CCP whereas 2 patients (4%) was negative for both (**Table – 1**).

Table - 1: No. of patients with various serology and inflammatory markets.

	No. of pts	%
RF positive	40	80
Anti-CCP positive	37	74
Both RF, anti-CCP positive	30	60
Both RF, anti-CCP negative	02	04
Raised ESR	42	84
Raised CRP	45	90

Table - 2: Mean value of RF, DAS28, Larsen score, CRP and ESR in Anti-CCP positive and Anti-CCP negative patients.

Mean value of	Anti-CCP	
	Positive	Negative
RF level	199	67
DAS28	4.74	3.18
Larsen score	21.9	14.1
CRP	38	22
ESR	34	28

Table - 3: Mean value of DAS28, Larsen score, CRP and ESR in RF positive and RF negative patients.

Mean value of	RF positive	RF negative
DAS28	4.9	3.5
Larsen score	25.7	13.2
CRP	38	20
ESR	44	24.2

The level of RF was significantly high in anti-CCP positive patients than in anti-CCP negative patients ($p < 0.01$) means significant correlation between RF and anti-CCP. Larsen score, DAS28 were higher in anti-CCP positive patients than in anti-CCP negative patients. There were early destructive changes in joints in anti-CCP positive patients than in anti-CCP negative patients indicates prognostic value of anti-CCP antibody (**Table – 2**).

On other hand Larsen score and DAS28 were significantly higher in RF positive patients than in RF negative patients ($p < 0.05$, $P < 0.001$ respectively) as per **Table – 3**.

Discussion

The critical strategy to prevent joint damage in RA is to initiate DMARD early in course of the disease. The ability to predict severe disease outcome is as important as a correct diagnosis. Anti-CCP antibody and RF are shown as important serological marker for RA diagnosis and as a possible prognostic marker for development of early erosive disease [13, 14]. A recent study showed that in patients with synovitis of three months duration, a combination of anti-CCP and RF has high specificities (97%) and PPV (83%) for development of persistent RA. Furthermore Anti-CCP have been incorporated in to newly proposed diagnostic criteria for RA and proved to be strongly associated with erosive arthritis.

In this study, anti-CCP and RF positivities and the duration of the disease were independent predictors of erosive development, and

combination of these antibodies had the highest for erosive damage.

Most studies agree that a positive RF is an important predictor for joint damage over the years of disease. Jansen, et al. [5] concluded that radiographic progression at one year was predicted by positive RF. Similar to previous studies, our results suggested the prognostic value of RF.

It appears that anti-CCP antibodies have prognostic relevance similar to RF. Vencovsky, et al. [9] found that anti-CCP positivity was better than RF at predicting progression of Larsen score over 2 years. Also, in prospective study of 242 patients with early RA followed up for three years, the anti-CCP antibody results correlated with RF, but were better than RF as predictor of more aggressive disease [15].

Finally this study showed that the presence of anti-CCP and RF antibodies was associated with high probability of erosive disease. The combined use of RF and anti-CCP had greater specificity and PPV for erosive damage than anti-CCP or RF alone.

Conclusion

- RA is more common in female.
- There is significant correlation between RF and anti-CCP antibody.
- Anti-CCP antibody and RF both in combination were associated with higher probabilities of erosive disease.

References

1. Forslind K, Ahlmen M, Eberhardt K, Hatstrom I, Svenson B. Prediction of radiological outcome in early RA in clinical practice. Role of antibodies to citrullinated peptide. *Ann Rheum Dis.*, 2004; 63: 1090-1095.
2. Raza K, Breese M, Nightingale P, Kumar K, potter T, Earlythers, et al. Predictive value of antibodies to cyclic citrullinated peptide in patients with

- early inflammatory arthritis. *J Rheumatol.*, 2005; 32: 231-238.
3. Vittecoq O, Pouplin S, Krzanowska K, Jousen –Beades F, Menard JF, Gaet A, et al. Rheumatoid factor is the strongest predictor of radiological progression of rheumatoid arthritis in three-year prospective study in community-recruited patients. *Rheumatology*, 2003; 42: 939-946.
 4. Goekoop-Ruiterman YP, de Vrus-Bouwstra JK, Allaart CF, van Zeben D, Kerstens PJ, et al. comparison of treatment strategies in early rheumatoid arthritis; a randomized trial. *Ann Intern Med.*, 2007; 146: 406-415.
 5. Jansen AL, van der Horst- Bruinsma I, van Schaardenburg D, Van de Stadt RJ, de Koning, Dijkmans BA. RF and anti-CCP differentiate RA from undifferentiated spondyloarthropathies in patients with early arthritists. *J Rheumatol.*, 2002 Oct; 29(10): 2074-6.
 6. Van Jaarveld CH, ter Borg EJ, Jacobs JW, Schellekens GA, Gmelig-Meylig FH, Van Booma-Frankfort C, et al. Prognostic value of the antiperinuclear factor and anti-CPP and RF in early RA. *Clin Exp Rheumatol.*, 1999; 17: 689-697.
 7. Lee DM, Schur PH. Clinical utility of Anti-ccp assay in patients with rheumatic disease. *Ann rheum Dis.*, 2003; 62: 870-874.
 8. Kastbam A, Strandberg G, Lindroos A, Skogh T. Anti ccp antibody test predict disease course during 3 year. In early RA (The Swedish TIRA Project). *Ann. Rheum Dis.*, 2004; 63: 1085-1084.
 9. Vencovsky J, Machacek S, Sedoval, Kafkova J, Gatterova J, Pesakova V, et al. Autoantibodies can ben prognostic markers of erosive disease in early RA. *Ann. Rheum Dis.*, 2003; 62: 424-30.
 10. Arhett FC, Edworthy SM, Bloch DA, Mcshane DJ, Fries JF, Cooper NS, et al. The American rheumatism association 1987 revised criteria for the classification of rheumatoid arthiristis. *Arthritis rheum.*, 1988; 31: 315-324.
 11. Larsen A, Dale K, Eek M. radiographic evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta radi-ol Diagn.*, 1977; 18: 481-491.
 12. Jansen LM, van der Horst – Bruinsma IE, van Schaardenburg D, Benzemer PD, Dijkmans BA. Predictors of radiographic joint damage in patients with early RA. *Ann Rheum Dis.*, 2001; 60: 924-927.
 13. Berglin E., Johansson T, Sundin U. Radiological out come in RA is predicted by presence of anti-CCP before and at disease onset. *Ann Rheum Dis.*, 2006; 65: 453-458.
 14. Avouac J, Gossec L, Dougados M. diagnostic and predictive value of anti-CCP in RA: a systemic literature review. *Ann Rheum Dis.*, 2006; 65: 845-851.
 15. Winska Wiloch H, Thompson K, Young A, Corbett M, Shipley M, Hay F. RF as marker of late erosive changes in RA. *Scand J Rheumatology*, 1988; 75: 238-243.